



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**Embryos of *Angiopteris* and *Kaulfussia*.**—In connection with a study of these two genera, CAMPBELL<sup>26</sup> has presented the embryo-formation of Marattiaceae. He thinks it probable that in all cases the stem, leaf, and root are epibasal in origin. In *Danaea* the primary hypobasal cell forms a suspensor, so that the foot also is epibasal; and in *Kaulfussia* and *Angiopteris* the foot is also partly epibasal in origin. The root arises endogenously as a secondary structure, and in its growth almost obliterates the foot, which is very large in the young embryo.—J. M. C.

**Anatomy of *Calamostachys*.**—HICKLING<sup>27</sup> has studied new sections of *C. Binneyana*, and has come to the conclusion that the so-called fertile or sporangiophore-bearing “nodes” are not nodes in the same sense as the bract-bearing nodes. He shows that the sporangiophore trace arises from the node that supplies the whorl of bracts below, and he believes this to be a general characteristic of calamitean strobili, the point of insertion of the sporangiophores on the internode being very variable.—J. M. C.

---

<sup>26</sup> CAMPBELL, D. H., The embryo and young sporophyte of *Angiopteris* and *Kaulfussia*. Ann. Jard. Bot. Buitenzorg II. Suppl. III. pp. 69–82. pls. 6, 7. 1909.

<sup>27</sup> HICKLING, GEORGE, The anatomy of *Calamostachys Binneyana* Schimper. Mem. and Proc. Manchester Lit. and Phil. Soc. 54: no. 17. pp. 16. pl. 1. 1910.